ISSN 2729-8043

INTERNATIONAL JOURNAL OF ORIGINAL MEDICINE

Original Medicine. Volume 3 Issue 2 2024
ISSN 2729-8043
https://www.original-medicine.eu/
Impact factor 8
Original Medicine is an open-access publication, providing authors with continuous publication of original research across a broad spectrum of medical, scientific disciplines and sub-specialties. The Original Medicine review process emphasizes submissions' scientific, technical and ethical validity. Novelty or potential for impact is not considered during the manuscript's evaluation or adjudication.

Manuscripts typed on our article template can be submitted through our website here. Alternatively, authors can send papers as an email attachment to info@original-medicine.eu

Original Medicine.
ISSN 2729-8043 https://www.original-medicine.eu
Michalská 2, Bratislava, Slovakia, 81121
E-mail: info@original-medicine.eu

Original Medicine. Volume 3 Issue 2 2024
ISSN 2729-8043
https://www.original-medicine.eu/
Impact factor 8

RELATIONSHIP BETWEEN METABOLIC SYNDROME AND ACQUIRED PREMATURE EJACULATION

Rahmonova Vasila Farkhodovna,
REPUBLICAN SPECIALIZED SCIENTIFIC
AND PRACTICAL MEDICAL CENTER OF ENDOCRINOLOGY.
NAMED AFTER ACADEMICIAN Y.KH. TURAKULOVA

Abstract: One of the leading global public-health burdens is metabolic syndrome (MetS), despite the many advances in pharmacotherapies. MetS, now known as “developmental origins of health and disease” (DOHaD), can have its origins in early life. Offspring MetS can be programmed by various adverse early-life conditions, such as nutrition imbalance, maternal conditions or diseases, maternal chemical exposure, and medication use. Conversely, early interventions have shown potential to revoke programming processes to prevent MetS of developmental origins, namely reprogramming.

Keywords: metabolic syndrome, premature ejaculation, developmental origins of health and disease (DOHaD)

A prospective and comparative investigation on the association between acquired premature ejaculation (PE) and metabolic syndrome would aim to explore potential links between these two health conditions. Premature ejaculation is a common sexual dysfunction, while metabolic syndrome is a cluster of risk factors that increase the risk of cardiovascular diseases, type 2 diabetes, and other health issues. This study would likely address several key points:

1. Study Design. Prospective Nature: A prospective study would involve following participants over time to observe the development of acquired PE in individuals with or without metabolic syndrome. Comparative Aspect: The study would compare individuals diagnosed with metabolic syndrome to those without, examining the prevalence or incidence of PE in both groups.

2. Study Population. The study would recruit a diverse population of men, both with and without metabolic syndrome, to assess whether there is a higher likelihood of developing acquired PE in those with metabolic syndrome. Consideration would be given to age, lifestyle factors (such as obesity, sedentary lifestyle, and diet), and other co-morbid conditions.

3. Variables and Data Collection. Primary Variables: Presence or absence of acquired PE and metabolic syndrome. Secondary Variables: Body mass index (BMI), waist circumference, blood pressure, lipid profile, fasting blood glucose, and other

factors related to metabolic health. Psychological and Lifestyle Factors: Assessment of stress, anxiety, and psychological factors that may 050% of men with MetS may experience PE.

2. Hormonal Imbalance and Vascular Factors: Testosterone Deficiency: Men with MetS often have lower levels of testosterone, which is linked to both erectile dysfunction and premature ejaculation. Testosterone plays a role in sexual function, and its decline can lead to faster ejaculation times and difficulty in regulating ejaculatory reflexes.

3. Insulin Resistance: Metabolic syndrome is marked by insulin resistance, which is believed to affect the nervous system and lead to increased excitability of the ejaculatory reflex.

Endothelial Dysfunction: Poor vascular health (a key feature of MetS) may reduce blood flow to the pelvic region, affecting the function of tissues involved in sexual performance. This might contribute to difficulties in controlling ejaculation.

3. Psychological Stress and PE: Men with metabolic syndrome often suffer from higher levels of stress, anxiety, and depression, which are known psychological factors contributing to PE. The chronic nature of MetS can add to mental health challenges, compounding sexual dysfunction issues like premature ejaculation.

4. Obesity and Physical Health: Obesity, a core component of metabolic syndrome, is associated with an increased risk of premature ejaculation. Increased fat mass is associated with chronic inflammation, hormonal disturbances, and poor cardiovascular health, all of which can contribute to both PE and erectile dysfunction.

5. Lipid Profiles and Ejaculatory Control: Abnormal cholesterol levels (particularly high triglycerides and low HDL) found in MetS patients mayRelationship between metabolic syndrome and acquired premature ejaculation have indirect effects on nervous system function, including those pathways that control ejaculation.

Implications for Clinical Practice: Management of MetS May Improve PE Symptoms: Some research suggests that addressing the underlying metabolic issues through lifestyle interventions (such as weight loss, physical exercise, and improved diet) may improve premature ejaculation symptoms. Medical management of blood sugar, cholesterol, and blood pressure could also have a beneficial impact on sexual health.

Pharmacological Treatments: For men with both MetS and PE, combining standard treatments for PE (such as selective serotonin reuptake inhibitors, SSRIs, or topical anesthetics) with medications for metabolic syndrome may offer comprehensive management.

The results of existing studies strongly suggest a positive association between early climax (PE) and metabolic syndrome. Men with metabolic syndrome are at greater risk for developing sexual dysfunctions, including PE, due to the complex interplay of hormonal imbalances, vascular issues, and psychological stress. Further research with larger sample sizes and longer follow-up periods is necessary to establish definitive causal pathways and to develop targeted therapies that address both metabolic and sexual health.

MetS may be an independent predisposing factor for the development of acquired PE. Therefore, MetS may be an independent predisposing factor for acquired PE, and the effective prevention and treatment of MetS could also be important for the prevention of acquired PE or ejaculatory dysfunction.

References

1. Alberti, K.G.; Eckel, R.H.; Grundy, S.M.; Zimmet, P.Z.; Cleeman, J.I.; Donato, K.A.; Fruchart, J.C.; James, W.P.; Loria, C.M.; Smith, S.C., Jr.; et al. Harmonizing the metabolic syndrome: A joint interim statement of the International Diabetes Federation Task Force on Epidemiology and Prevention; National Heart, Lung, and Blood Institute; American Heart Association; World Heart Federation; International Atherosclerosis Society; and International Association for the Study of Obesity. Circulation 2009, 120,1640–1645.

2. Parnham A, Serefoglu EC. Classification and definition of premature ejaculation. Transl Androl Urol 2016; 5:416–423

3. Kaur J. A comprehensive review on metabolic syndrome. Cardiol Res Pract 2014; 2014:943162.

4. The relationship between acquired premature ejaculation and metabolic syndrome: a prospective, comparative study. D Bolat et al. Int J Impot Res. 2017 May.

5. Metabolic Syndrome Is an Independent Risk Factor for Acquired Premature Ejaculation. Jeh SU, et al. World J Mens Health. 2019. PMID: 30588783 Free PMC article.

6. https://pubmed.ncbi.nlm.nih.gov/33831976/